



INVESTOR IN PEOPLE

6804/2702

## PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

The Patent Office  
Concept House  
Cardiff Road  
Newport

South Wales  
NP10 8QQ

REC'D 30 JUL 2004

WIPO


PCT

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed

*P. Mahoney*

Dated

6 July 2004


# BEST AVAILABLE COPY

23. JUN. 2003 17:23

HARRISON GODDARD

NO. 770

Patents Form 1/77

Patents Act 1977  
(Rule 16)The  
Patent  
Office24JUN03 E817204-1 002973  
P01/7700 0.00-0314624.8

The Patent Office

Cardiff Road  
Newport  
South Wales  
NP8 1RH**Request for grant of a patent**

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

**1. Your reference**

P102877GB

**2. Patent application number**

(The Patent Office will fill in this part)

0314624.8

**3. Full name, address and postcode of the or of each applicant (underline all surnames)**Advanced Bionutrition(Europe) Limited  
15 Grafton Road  
Worthing  
West Sussex  
BN11 1QR  
GB

Patents ADP number (if you know it)

If the applicant is a corporate body, give the country/state of its incorporation

8658577001  
GB**4. Title of the invention**

Inflammatory Disease Treatment

**5. Name of your agent (if you have one)**

Harrison Goddard Foote

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

31 St Saviourgate  
YORK  
YO1 8NQ

Patents ADP number (if you know it)

07914237002 ✓

**6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number**

Country

Priority application number  
(if you know it)Date of filing  
(day / month / year)**7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application**

Number of earlier application

Date of filing  
(day / month / year)**8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:**

- a) any applicant named in part 3 is not an inventor, or  
 b) there is an inventor who is not named as an applicant, or  
 c) any named applicant is a corporate body.  
 See note (d))
- yes

Patents Form 1/77

0072563 23 Jun 03 05:28

**Patents Form 1/77**

Enter the number of sheets for any of the following items you are filing with this form.  
Do not count copies of the same document

**Continuation sheets of this form**

Description	16	/
Claim(s)	4	/
Abstract	1	/ <i>mw</i>
Drawing(s)	-	

10. If you are also filing any of the following, state how many against each item.

**Priority documents****Translations of priority documents**

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patents Form 9/77) 1 / *mw*

Request for substantive examination (Patents Form 10/77)

Any other documents (please specify)

11.

I/We request the grant of a patent on the basis of this application.

Signature

Date

23.6.03

12. Name and daytime telephone number of person to contact in the United Kingdom

Dr Rob Docherty

01904 732120

**Warning**

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

**Notes**

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

**Patents Form 1/77**

### INFLAMMATORY DISEASE TREATMENT

5 The invention relates to a composition comprising a source of long chain polyunsaturated fatty acids, for example, docosahexaenoic acid (DHA), and a carotenoid, for example, astaxanthin, and other nutrients for prophylactic and/or therapeutic use in the healing of trauma- and stress-induced inflammatory conditions.

#### Background to the invention

10 Inflammation is a complex stereotypical reaction of the body responding to damage of its cells and vascularised tissues. The damaged sites are susceptible to infiltration by a multitude of pathogens including viruses, bacteria, fungi, and protozoan and metazoan parasites, as well as cancerous cells and other harmful agents and so the animal defends itself by initiating an inflammatory reaction at the damaged site.

15 The inflammatory reaction is phylogenetically and ontogenetically the oldest defence mechanism and both the innate and adaptive immune systems in vertebrates are triggered to destroy the infectious agent(s). When a tissue has been traumatised, for example, by injury or surgery, and is thus susceptible to infection, three key steps in the inflammatory response are initiated; (1) vasodilation, which enables an increased  
20 blood supply to the traumatised tissue; (2), increased capillary permeability caused by retraction of the endothelial cells allowing soluble mediators of immunity to reach the site of inflammation; and (3) migration of leukocytes (neutrophils; monocytes and lymphocytes) out of the capillaries into the surrounding tissues.

25 The development of inflammatory reactions is controlled in part by pro-inflammatory cytokines (e.g. interleukin-1, tumour necrosis factor alpha); by lipid mediators released from different cells (e.g. prostaglandin's and leukotrienes); by cell-derived vasoactive mediators released from mast cells, basophils and platelets (e.g. arachidonic acid metabolites; platelet activating factors amines: serotonin, histamine; endothelins) and by plasma-derived vasoactive mediators (e.g. kinins and components  
30 of the complement, coagulation and fibrinolytic cascades).

Chronic inflammation is an inflammatory response of prolonged duration - weeks, months, or even indefinitely - whose extended time course is provoked by the persistence of the causative stimulus to inflammation within the tissue. The inflammatory process inevitably causes tissue damage and is accompanied by mis-  
5 directed attempts at simultaneous healing and repair. The exact nature, extent and time course of chronic inflammation is variable, and depends on a balance between the causative agent and the attempts of the body to remove it.

Chronic inflammation may develop either as a progression from acute inflammation if  
10 the original stimulus persists, or after repeated episodes of acute inflammation or *de novo* if the causative agent produces only a mild acute response.

Aetiological agents producing chronic inflammation include, but are not limited to: infectious organisms that can avoid or resist host defences and so persist in the tissue for a prolonged period; infectious organisms that are not innately resistant but persist  
15 in damaged regions where they are protected from host defences; irritant non-living foreign material that cannot be removed by enzymatic breakdown or phagocytosis; or where the stimuli is a "normal" tissue component, causing an auto-immune disease.

There is a vast array of diseases exhibiting a chronic inflammatory component. These include but are not limited to: inflammatory joint diseases (e.g., rheumatoid arthritis,  
20 osteoarthritis, polyarthritis and gout), chronic inflammatory connective tissue diseases (e.g., lupus erythematosus, scleroderma, Sjorgen's syndrome, poly- and dermatomyositis, vasculitis, mixed connective tissue disease (MCTD), tendonitis, synovitis, bacterial endocarditis, osteomyelitis and psoriasis), chronic inflammatory lung diseases (e.g., chronic respiratory disease, pneumonia, fibrosing alveolitis,  
25 chronic bronchitis, bronchiectasis, emphysema, silicosis and other pneumoconiosis and tuberculosis), chronic inflammatory bowel and gastro-intestinal tract inflammatory diseases (e.g., ulcerative colitis and Crohn's disease), chronic neural inflammatory diseases (e.g., chronic inflammatory demyelinating polyradiculoneuropathy, chronic inflammatory demyelinating polyneuropathy,  
30 multiple sclerosis, Guillan-Barre Syndrome and myasthenia gravis), other inflammatory diseases (e.g., mastitis, laminitis, laryngitis, chronic cholecystitis, Hashimoto's thyroiditis, inflammatory breast disease); chronic inflammation caused

by an implanted foreign body in a wound; and acute inflammatory tissue damage due to muscle damage after eccentric exercise (e.g., delayed onset muscle soreness - DOMS).

The usual mode of treatment for chronic inflammatory conditions is by administration of non-steroidal anti-inflammatory drugs (NSAID's) such as Diclofenac, Ibuprofen, Aspirin, Phenylbutazone, Indomethacin, Naproxen and Piroxicam. Although NSAID's can be effective, they are known to be associated with a number of side effects and adverse reactions. These may include gastro-intestinal problems such as dyspepsia, ulceration and haemorrhage, sleepiness, nausea or vomiting, constipation, allergic reactions and occasionally hepatotoxicity. Frequently the margin between effective dose and toxic dose is quite small (i.e., 2-3 -fold). In spite of these side effects, the use of NSAID's as anti-inflammatory agents is standard practice in human medicine and veterinary medicine. However, within veterinary medicine there is an increasing concern about their use in food animals because of the potential for accumulation of drugs such as phenylbutazone within the food chain.

It is the purpose of this invention to provide a natural alternative to anti-inflammatory drugs widely used to treat chronic inflammatory conditions of terrestrial animals including humans. The use of such an alternative will be safe and without side-effects or risks to the environment.

DHA is an omega-3 fatty acid and is the most abundant long chain polyunsaturated fatty acid (PUFA) in the grey matter of the brain and other neurological tissues. Omega-3 PUFAs, particularly eicosapentaenoic acid (EPA) are known to be beneficial in reducing incidence of coronary heart disease (Lands, Fish and Human Health 1986 Academic Press). The anti-inflammatory properties of omega-3 PUFAs are thought to be provided by their ability to replace arachidonic (ARA) acid in immune cells membranes. ARA, an omega-6 PUFA with 20 carbon atoms and 4 double bonds (C20:4), is the biochemical precursor for the production of 2-series prostaglandins and 4-series leukotrienes associated with a range of pro-inflammatory molecules and mediators and can therefore impact pathogenesis of inflammatory diseases. (P. Calder "n-3 Fatty Acids & Health Conference (December 1999) British Nutrition Foundation). EPA, an omega-3 PUFA with 20 carbon atoms and 5 double

bonds (C20:5) is the biochemical precursor for the production of 3-series prostaglandins and 2-series leukotrienes which are anti-inflammatory molecules. Thus, the balance of EPA and ARA is thought to significantly affect the balance of pro- and anti-inflammatory eicosanoid mediators. DHA, an omega-3 PUFA with 22 carbon atoms and 6 double bonds (C22:6) does not form eicosanoids (i.e., 20C prostaglandins or leukotrienes). Omega-3 fatty acids have a long history of use in animal feeding via use of cod liver oil, linseed and flax oil.

A metabolic pathway exists in mammals for the biosynthesis of DHA, but this is bio-energetically unfavourable (Crawford, P. AOCS, Short Course in Polyunsaturated Fatty Acids and Eicosanoids, pp270-295 (1987). The metabolism of omega-3 fatty acids is not well understood, thus precise clinical dosage rates and efficacy remain unknown. Mammals are thought to obtain most of their DHA from dietary sources.

Omega-3 and omega-6 fatty acids are found in cold-water marine fish; and fish oils are the primary commercial source of these fatty acids. Environmental pollution of fish introduces toxic factors such as dioxins and PCB's to the oils recovered from fish, which if ingested may adversely affect the health of all animals and may remain as residues in food animals rendering them problematic for human consumption.

Marine microorganisms are known to contain DHA, in particular dinoflagellates (Harrington et al "The Polyunsaturated Fatty Acids of Marine Dinoflagellates" J. Protozoal, 17:213-219 (1970)). Successful cultivation of these in commercial conditions is achievable (U.S 5,407,957). In adequate presence of Vitamin E up to animals can consume up to 2% of their diet as DHA when using fish oil, but higher levels result in malodorous products.

Astaxanthin is a carotenoid known to be partially degraded in the gastro-intestinal tract by oxidation. The presence of vitamins A, C, selenium, manganese, zinc and copper are known to alleviate this effect. Certain microorganisms including but not limited to algae and yeast are known to be prolific producers of astaxanthin. Both forms of algae, and yeast contain adequate combinations of the above elements to counteract the oxidative effect of digestive oxidation to both the lipids and the astaxanthin therein.

Other marine organisms, including but not limited to zooplankton, crustaceans, molluscs, and vertebrates, are also known to contain high levels of the carotenoid astaxanthin. It has been shown that in fish and crustaceans, astaxanthin is essential for growth and plays a vitamin-like role. Astaxanthin also appears to have some beneficial effects on mammals. Astaxanthin is an active ingredient in several patented medications for mammals. In an anti-stress formulation, it is claimed to enhance the effect of anti-stress agents administered to farm animals and household pets to minimise weight loss and reduced immunity due to crowding, extreme temperatures and other sudden environmental changes (U.S 5,937,790).

Esterified astaxanthin from the alga *Haematococcus pluvialis* is the preferred form in several oral prophylactic and therapeutic formulations for muscular dysfunction such as exertional rhabdomyolysis (also known as exertional myopathy, tying-up syndrome, azoturia, or Monday morning sickness) in horses (WO 99/11251), as well as for mastitis (mammary inflammation) in dairy cows (WO 98/30701), and for mammalian gastrointestinal tract inflammation due to infections by *Helicobacter* sp. bacteria (WO 98/37874).

The use of yeast in animal feed has a long and well-documented history. Recent changes to European law (EC Directive 87/153/EEC and associated reports) specifically in respect to the ability of a gastro-intestinal tract to resist overgrowth by any one component or strain is now active. Whilst the mode of action is not documented, it is thought that there are similarities between the action of the rumen and the cecal fermentation of mammals that rely on bacterial fermentation resulting in production of lactic acid. (Martin, S.A. and Nisbet, D.J., "Effect of direct-fed microbials on rumen microbial fermentation" J. Dairy Sci. 75:1736 (1992))

Recent research and meta-analysis shows yeast to improve digestion and availability of nutrients when nutritional demands are high. Positive effects on the efficacy of immune systems to increase macrophage activity against *E. coli* and *Salmonella typhitrium* have been shown. (Hatcher G. E., Lambretch R.S., "Augmentation of macrophage phagocytic activity by cell-free extracts of selected lactic acid producing bacteria" J. Dairy Sci. 76:2485 (1993) and Schiffrin, E.J. et al, "Immunomodulation



of human blood cells following the ingestion of lactic acid bacteria J. Dairy Sci 78:491 (1995))

5 An object of the present invention is to provide a dietary supplement to an animal, including humans, that will provide a protective benefit against inflammation (particularly to animals in high stress environments such as, but not limited to competition or transportation), and/or to be used therapeutically to further enhance the healing of trauma and stress-induced inflammatory conditions.

10 A further objective of this invention is to provide a natural alternative to anti-inflammatory drugs currently used in traditional veterinary and human medicine.

We have found that a combination of an omega-3 PUFA and astaxanthin provides an unexpectedly beneficial effect in the prevention of inflammation and further that these  
15 materials can be provided to an animal, including humans, in a natural and bioavailable form.

#### Summary of the Invention

20 According to an aspect of the invention there is provided a composition comprising at least one long chain polyunsaturated fatty acid and at least one carotenoid.

In a preferred embodiment of the invention said long chain fatty acid is a free fatty acid, or an ester thereof.

25 In a further preferred embodiment of the invention said long chain fatty acid is selected from the group consisting of: a triglyceride, diglyceride, monoglyceride, phospholipids, glycolipid, sphingolipid or sulpholipid.

In a further preferred embodiment of the invention said long chain fatty acid is  
30 docosahexaenoic acid.

In a preferred embodiment of the invention said docosahexaenoic acid is provided as an edible algae. Preferably said edible algae is selected from the group consisting of, but not limited to: *Crypthecodinium*; *Phaedactylum*; *Isochrysis*; *Schizochytrium*; *Thaustochytrium*; or *Ulkenia*.

- 5 In a preferred embodiment of the invention said long chain fatty acid is eicosapentaenoic acid.

In a preferred embodiment of the invention said eicosapentaenoic acid is provided as an edible algae. Preferably said edible algae is selected from the group consisting of, but not limited to: *Isochrysis*; *Nannochloris*; *Cyclotella*; *Phaeodactylum*, or *Navicula*,

- 10 In a yet further preferred embodiment of the invention said carotenoid is astaxanthin.

In a preferred embodiment of the invention said composition further comprises yeast.

In a further preferred embodiment of the invention said composition further comprises a further anti-inflammatory or antioxidant agent.

- 15 In a preferred embodiment of the invention said further anti-inflammatory or antioxidant agent is selected from the group consisting of, but not limited to: vitamin C, vitamin E, lycopene,  $\beta$ -carotene, lutein, organic selenium,  $\alpha$ -lipoic acid, glycine, taurine, methylsulfonylmethane, glutamine, arginine, cysteine, methionine, S-adenosylmethionine, nucleotides, nucleic acids, curcumin, green tea extract, green-lipped mussel extract (*Perna canaliculus*) or standardised herbal extracts such as  
20 *Phyllanthus amarus*, *Fructus Schisandra*, Chamomile, Blackcurrant leaf or Devil's claw.

According to a further aspect of the invention there is provided a composition according to any previous aspect or embodiment for use as a nutraceutical.

- 25 When administered, compositions of the present invention are administered in physiologically acceptable preparations. Such preparations may routinely contain physiologically acceptable concentrations of salt, buffering agents, preservatives, compatible carriers and optionally other therapeutic agents.

- 30 The compositions of the invention can be administered by any conventional route, including, but not limited to, injection, or gradual infusion over time. The

administration may, for example, be oral, intravenous, intraperitoneal, intramuscular, intracavity, subcutaneous, or transdermal. Preferably said compositions are administered orally in the feed or as a feed supplement. Alternatively, the compositions can be provided in the water or as a tonic.

5

The compositions of the invention are administered in effective amounts. An "effective amount" is that amount of a composition that alone, or together with further doses, produces the desired response. In the case of treating a particular disease, such as arthritis, the desired response is inhibiting the progression of the disease. This may  
10 involve only slowing the progression of the disease temporarily, although more preferably, it involves halting the progression of the disease permanently. This can be monitored by routine methods.

Such amounts will depend, of course, on the particular condition being treated, the  
15 severity of the condition, the individual patient parameters including age, physical condition, size and weight, the duration of the treatment, the nature of concurrent therapy (if any), the specific route of administration and like factors within the knowledge and expertise of the health practitioner. These factors are well known to those of ordinary skill in the art and can be addressed with no more than routine  
20 experimentation. It is generally preferred that a maximum dose of the individual components or combinations thereof be used, that is, the highest safe dose according to sound medical judgment. The compositions used in the foregoing methods preferably are sterile and contain an effective amount of the active agents for producing the desired response in a unit of weight or volume suitable for  
25 administration to a patient.

In general, the active agent DHA is formulated and administered in doses between 0.05-500mg/kg body weight, preferably between 0.5-15 mg/kg body weight, and most preferably between 1-3mg /kg body weight. The active agent astaxanthin is  
30 formulated and administered in doses between 0.0005mg-5mg/kg body weight, preferably between 0.0015-0.15mg/kg body weight, and most preferably between 0.0075-0.0225mg/kg body weight according to any standard procedure in the art.

Compositions may be combined, if desired, with a physiologically-acceptable carrier. The term "physiologically-acceptable carrier" as used herein means one or more compatible solid or liquid fillers, diluents or encapsulating substances which are suitable for administration into a human or animal. The term "carrier" denotes an  
5 organic or inorganic ingredient, natural or synthetic, with which the active ingredient is combined to facilitate the application. The compositions may contain suitable buffering agents.

The compositions may conveniently be presented in unit dosage form and may be  
10 prepared by any of the methods well-known in the art of pharmacy. All methods include the step of bringing the active agent into association with a carrier which constitutes one or more accessory ingredients. In general, the compositions are prepared by uniformly and intimately bringing the active compound into association with a liquid carrier, a finely divided solid carrier, or both, and then, if necessary,  
15 shaping the product. This can then be either administered directly to the animal or added to food.

Compositions suitable for oral administration may be presented as discrete units, such as capsules, tablets, lozenges, each containing a predetermined amount of the active  
20 agent. Other compositions include suspensions in aqueous liquids or non-aqueous liquids such as a syrup, elixir, tonic, or an emulsion.

According to a further aspect of the invention there is provided the use of a composition according to the invention for the manufacture of a medicament for use  
25 in the treatment of inflammatory conditions.

In a preferred embodiment of the invention said inflammatory condition is selected from the group consisting of, but not limited to: inflammatory joint diseases (e.g. rheumatoid arthritis, osteoarthritis, polyarthritis and gout); chronic inflammatory connective tissue diseases (e.g. lupus erythematosus, scleroderma, Sjorgen's  
30 syndrome, poly- and dermatomyositis, vasculitis ); mixed connective tissue disease (MCTD) (e.g. tendonitis, synovitis, bacterial endocarditis, osteomyelitis and psoriasis ); chronic inflammatory lung diseases (e.g. chronic respiratory disease, pneumonia, fibrosing alveolitis, chronic bronchitis, bronchiectasis, emphysema,

silicosis and other pneumoconiosis and tuberculosis); chronic inflammatory bowel and gastro-intestinal tract inflammatory diseases (e.g. ulcerative colitis and Crohn's disease); chronic neural inflammatory diseases (e.g. chronic inflammatory demyelinating polyradiculoneuropathy, chronic inflammatory demyelinating  
5 polyneuropathy, multiple sclerosis, Guillan-Barre Syndrome and myasthenia gravis); and other inflammatory diseases including, mastitis, laminitis, laryngitis, chronic cholecystitis, Hashimoto's thyroiditis, inflammatory breast disease; chronic inflammation caused by an implanted foreign body in a wound; and acute inflammatory tissue damage due to muscle damage after eccentric exercise (e.g.,  
10 delayed onset muscle soreness - DOMS).

According to a further aspect of the invention there is provided a food stuff wherein said food stuff comprises a composition according to any previous aspect of embodiment.

According to a further aspect of the invention there is provided a method to treat an  
15 animal suffering from an inflammatory condition or disease comprising administering to said animal an effective amount of at least one long chain polyunsaturated fatty acid and at least one carotenoid.

In a preferred method of the invention said long chain fatty acid is a free fatty acid, or  
20 an ester thereof.

In a further preferred method of the invention said long chain fatty acid is selected from the group consisting of: a triglyceride, diglyceride, monoglyceride, phospholipids, glycolipid, sphingolipid or sulpholipid.  
25

In a further preferred method of the invention said long chain fatty acid is docosahexanoic acid.

In a yet further preferred method of the invention said carotenoid is astaxanthin.

In a further preferred method of the invention said animal is administered a further  
30 anti-inflammatory agent.

In a preferred method of the invention said disease or condition is selected from the group consisting of but not limited to: inflammatory joint diseases (e.g. rheumatoid arthritis, osteoarthritis, polyarthritis and gout); chronic inflammatory connective tissue diseases (e.g. lupus erythematosus, scleroderma, Sjorgen's syndrome, poly- and dermatomyositis, vasculitis); mixed connective tissue disease (MCTD)(e.g. tendonitis, synovitis, bacterial endocarditis, osteomyelitis and psoriasis); chronic inflammatory lung diseases (e.g. chronic respiratory disease, pneumonia, fibrosing alveolitis, chronic bronchitis, bronchiectasis, emphysema, silicosis and other pneumoconiosis and tuberculosis); chronic inflammatory bowel and gastro-intestinal tract inflammatory diseases (e.g. ulcerative colitis and Crohn's disease); chronic neural inflammatory diseases (e.g. chronic inflammatory demyelinating polyradiculoneuropathy, chronic inflammatory demyelinating polyneuropathy, multiple sclerosis, Guillan-Barre Syndrome and myasthenia gravis); and other inflammatory diseases including, mastitis, laminitis, laryngitis, chronic cholecystitis, Hashimoto's thyroiditis, inflammatory breast disease; chronic inflammation caused by an implanted foreign body in a wound; and acute inflammatory tissue damage due to muscle damage after eccentric exercise (e.g., delayed onset muscle soreness - DOMS).

In a preferred method of the invention said animal is a terrestrial animal

20 In a further preferred method of the invention said animal is a companion or performance animal.

In a further preferred method of the invention said animal is selected from the group consisting of: human, horse, cow; sheep; goat; mink; pig; dog; cat; hamster; mouse; 25 rabbit; pot bellied pig; rat, gerbil, guinea pig.

In a preferred method of the invention said animal is a horse.

In a further preferred method of the invention said animal is a human.

30

An embodiment of the invention will now be described by example only and with reference to the following materials, methods and examples.

## Materials and Methods

### Sources of DHA and astaxanthin:

DHA can be found in oils extracted from marine animals and organisms, including algae. Suitable commercial sources of DHA include, but are not limited to algae such as *Crypthecodinium*; *Phaedactylum*; *Isochrysis*; *Schizochytrium*; *Thaustochytrium*; or *Ulkenia*, or purified, or semipurified lipid products from these species.

Alternatively DHA can be provided by commercially available marine oils which typically contain levels between 15% and 25% DHA and between 5% and 15% EPA (w/w). Suitable marine oils include, but are not limited to: crude or processed fish oil, krill oil, squid oil, or refining or processing coproducts from the manufacture of these oils.

Suitable sources of commercially available astaxanthin include, but are not limited to: the dried algae product *Haematococcus pluvialis* (Cyanotech Corp, USA), dehydrated yeast product *Phaffia rhodozyma* (Igene Corp, USA). Alternatively the commercially available synthetic form of astaxanthin may be used (Roche, Switzerland; BASF, Germany).

**TABLE 1. EXAMPLES OF SUITABLE PRODUCT COMPOSITIONS**

#### Formula 1.

Ingredient	%
<i>Crypthecodinium cohnii</i> (dried biomass)	60
<i>Haematococcus</i> (dried biomass)	5
Inactivated Brewers dried yeast	35

#### Formula 2.

Ingredient	%
<i>Crypthecodinium cohnii</i> (dried biomass)	40.0
<i>Haematococcus</i> (dried biomass)	3.4
Inactivated Brewers dried yeast	23.3
Grass Meal	33.3

**Formula 3.**

Ingredient	%
Schizochytrium sp. (dried biomass)	60
Haematococcus (dried biomass)	5
Inactivated Brewers dried yeast	35

**Formula 4.**

Ingredient	%
Schizochytrium sp. (dried biomass)	40.0
Haematococcus (dried biomass)	3.4
Inactivated Brewers dried yeast	23.3
Grass Meal	33.3

5

**Formula 5.**

Ingredient	%
Fish oil (microencapsulated)	75
Haematococcus (dried biomass)	3
Inactivated Brewers dried yeast	10
Grass Meal	13

**Formula 6**

Ingredient	%
DHA algal oil (DHASCO)®	15
Astaxanthin (synthetic)	0.045
$\alpha$ -Lipoic acid	0.2
Natural flavours	0.3
Maltodextrin 20 DE	to 100

10



**Formula 7**

Ingredient	%
DHA algal oil (DHASCO)®	15
Astaxanthin (Pfaffia)	0.045
Methyl sulfonyl methane (organic)	0.2
Natural flavours	0.3
Maltodextrin 20DE	to 100

**EXAMPLE 1**

5

**Case Study 1**

A 26-year-old, 15.2hh  $\frac{3}{4}$  thoroughbred gelding had undergone metacarpal surgery on his off-fore knee 15 months prior to intervention. The animal was suffering osteoarthritis, general stiffness more deterioration in ability to movement when cold.

- 10 Long-term soft tissue swelling at the site of the injury and surgery remained round knee joint decreasing possibility of flexion further. Oedema surrounding ligaments and tendons occurred upon inactivity (e.g. stabling)

- 15 Dietary intake had included a range of herbal supplements, but no chemical phenylbutazone or other anti-inflammatory substances; 95% forage based with 1kg per day cereal mix. Initial dose of 0.5g formula 1 per day was included in the diet, increasing over 14 days to 8g intervention per day. Soft tissue swelling reduced above and below knee within 7 days (5mm decrease in circumference above knee and 4mm below knee), oedema reduced in hind lower limbs, stiffness on activity following rest
- 20 was noticeably reduced. After 21 days general alertness and overall health was noticed to have improved, e.g. coat condition. Horse was more eager to canter in field and less prone to stumbling on off fore.

25

**EXAMPLE 2****Case Study 2**

The subject was a 23-year-old, 13.2hh cob mare exhibiting old age related stiffness, notably following work on hard ground and when weather was cold and wet. Clinical symptoms were reluctance to move quickly whether ridden or in-hand, stiffness when moving out of stable following period of inactivity, oedema in lower limbs, slight grumpy manner when being handled and reluctance to engage in spontaneous movement in field. General health was good. Diet was 95% forage based, with small amount of soaked sugar beet pulp per day. No drug therapy was used, but pony has previously received phenylbutazone for stiffness and swellings.

Initial dose of 5g formula 1 per day for 5 days showed dramatic improvement with eagerness to move both ridden, in-hand and when free. Energy levels increased with improvement in disposition when handled, pony started to jump out of field over 1 metre 10cm high fence which previously was impossible for her, stride length of hind limbs increased to allow hoof prints of front hooves to be covered by hind hooves. Dose reduced to 2.5g per day, improvements still noticeable and oedema in lower limbs reduced. Maintenance dose of 2g per day showed no loss of activity. Comments on improvement in action, attitude and ability of pony noted by owner, farrier and instructor, none of whom were aware of the dietary changes made.

**EXAMPLE 3****Case Study 3**

Subject was a 11-year-old 12hh show pony suffering from laminitis, and not ridden due to chronic lameness, possibly due to muscle, shoulder injury sustained 22 months prior to test. No improvement when given phenylbutazone or other anti-inflammatory substances, difficult for farrier to work on as pony unable to move leg away from body at an angle; could not hold balance with foot off ground and was very stiff and sore for 2 to 5 days following attention from farrier, a result of the injury. 95% forage based diet with 1kg cereal inclusion per day, turn out in field but no exercise. Pony very stiff at all times, movement across uneven terrain difficult, not able to be ridden.

Initial dose of 2.5g formula 1 per day for 5 days showed dramatic improvement with pony much freer in action. Pony jumped out of field over 1meter 10cm fencing, landing on hard ground and was still sound, even the next day. After 4 weeks farrier shod pony and used him as an example to train other farriers because of his history of laminitis. Pony's ability to move leg, shoulder and withstand repeated lifting of legs was totally unexpected. Maintenance dose of 2g per day was used thereafter.

#### EXAMPLE 4

##### 10 Case Study 4

Subject was a 10-year-old King Charles Cavalier spaniel diagnosed with rheumatism in left hip, noticeable stiffness and inability to use hind limbs after exercise and worse in cold weather. Daily treatments with 0.5 g of Formula 1 resulted in improvement in coat and ability to exercise without pain (lifting of leg, limping, stopping suddenly) within 4 days. Dose was dropped to 0.5g per day on alternate days after 10 days of initial treatment and the animal continued to improve.

#### EXAMPLE 5

##### 20 Case Study 5

Subject was an 8 year old miniature Poodle, with general stiffness, unwillingness to jump on chairs, not using one hind limb when walking, notably stiff when getting up after rest, difficulty in using stairs of house and not playing with other dogs.

25 Intervention with Formula \_\_\_ at a dose of 0.25g per day with food for 7 days showed marked improvement in dogs movement, ability to jump on chairs/laps, and speed of ascent and descent of stairs. Play with other dog was initiated and speed of game was increased. Level of dose was maintained with overall improvement in dog's quality of life seen and overall condition.

30

## CLAIMS

1. A nutraceutical for use in the treatment of inflammatory conditions comprising docosahexaenoic acid or an ester thereof and at least one carotenoid.
2. The nutraceutical according to Claim 1 wherein the ester of docosahexaenoic acid is selected from the group consisting of: a triglyceride, diglyceride, monoglyceride, phospholipids, glycolipid, sphingolipid or sulpholipid.
3. The nutraceutical of Claims 1 or 2 wherein said docosahexaenoic acid is provided as an algae or fraction thereof.
4. The nutraceutical of Claim 3 wherein said algae is selected from the group consisting of: *Cryptocodinium*; *Phaedactylum*; *Isochrysis*; *Schizochytrium*; *Thaustochytrium*; or *Ulkenia*.
5. The nutraceutical of Claims 1 or 2 wherein said docosahexaenoic acid is provided as a fish oil or fraction thereof.
6. The nutraceutical according to any of Claims 1-5 wherein said carotenoid is selected from the group consisting of astaxanthin, zeaxanthin, lycopene, lutein, or carotene.
7. The nutraceutical according to any of Claims 1-6 wherein said carotenoid is from a microbial source.
8. The nutraceutical according to Claim 7 wherein said microbe is *Haematococcus* or *Pfaffia*.
9. The nutraceutical according to any of Claims 1-8 wherein said docosahexaenoic acid is provided to an animal at a dose of between 0.05 and 500 mg/kg body weight and said carotenoid is provided at a dose of between 0.5 and 5,000 ug/kg body weight.
10. The nutraceutical according to any of Claims 1-8 wherein said docosahexaenoic acid is provided to an animal at a dose of between 0.5 and 15 mg/kg body weight and said carotenoid is provided at a dose of between 1.5 and 150 ug/kg body weight.

11. The nutraceutical according to any of Claims 1-8 wherein said docosahexaenoic acid is provided to an animal at a dose of between 1 and 3 mg/kg body weight and said carotenoid is provided at a dose of between 7.5 and 22.5 ug/kg body weight.
- 5 12. The nutraceutical according to any of Claims 1-11 wherein said composition further comprises yeast.
13. The nutraceutical according to any of Claims 1-12 wherein said composition further comprises an anti-inflammatory agent.
- 10 14. The nutraceutical according to Claim 13 wherein said further anti-inflammatory agent is selected from the group consisting of: vitamin C, vitamin E, lycopene,  $\beta$ -carotene, lutein, organic selenium,  $\alpha$ -lipoic acid, methylsulfonylmethane, glutathione, taurine, glycine, glutamine, arginine, cysteine, methionine, S-adenosylmethionine, nucleotides, nucleic acids, curcumin, green tea extract, green-lipped mussel extract (*Perna canaliculus*), or standardised herbal extracts such as *Phyllanthus amarus*, *Fructus Schisandra*, Chamomile, Blackcurrant leaf, Devil's claw.
- 15 15. The nutraceutical according to any of Claims 1-14 wherein said inflammatory condition is selected from the group consisting of: inflammatory joint diseases e.g. rheumatoid arthritis, osteoarthritis, polyarthritis and gout; chronic inflammatory connective tissue diseases e.g. lupus erythematosus, scleroderma, Sjorgen's syndrome, poly- and dermatomyositis, vasculitis; mixed connective tissue disease (MCTD), e.g. tendonitis, synovitis, bacterial endocarditis, osteomyelitis and psoriasis; chronic inflammatory lung diseases e.g. chronic respiratory disease, pneumonia, fibrosing alveolitis, chronic bronchitis, bronchiectasis, emphysema, silicosis and other pneumoconiosis and tuberculosis; chronic inflammatory bowel and gastro-intestinal tract inflammatory diseases e.g. ulcerative colitis and Crohn's disease; chronic neural inflammatory diseases e.g. chronic inflammatory demyelinating polyradiculoneuropathy, chronic inflammatory demyelinating polyneuropathy, multiple sclerosis, Guillan-Barre Syndrome and myasthenia gravis; and including, mastitis, laminitis, laryngitis, chronic cholecystitis, Hashimoto's thyroiditis, inflammatory breast disease, and chronic inflammation.
- 20 25 30

16. A composition comprising and algal source of docosahexaenoic acid, a microbial source of astaxanthin, and grass meal.
17. The composition of Claim 16 wherein the algae is selected from the group consisting of: *Cryptocodinium*; *Phaedactylum*; *Isochrysis*; *Schizochytrium*; *Thaustochytrium*; or *Ulkenia*,
18. The composition of Claim 16 or 17 wherein the microbial source of astaxanthin is *Haematococcus* or *Pfaffia*.
19. The composition of Claims 16-18 further comprising inactivated brewers yeast.
20. The composition of any of claims 16-19 further comprising an anti-inflammatory agent selected from the group consisting of: vitamin C, vitamin E, lycopene,  $\beta$ -carotene, lutein, organic selenium,  $\alpha$ -lipoic acid, methylsulfonylmethane, glutathione, taurine, glycine, glutamine, arginine, cysteine, methionine, S-adenosylmethionine, nucleotides, nucleic acids, curcumin, green tea extract, green-lipped mussel extract (*Perna canaliculus*), or standardised herbal extracts such as *Phyllanthus amarus*, *Fructus Schisandra*, *Chamomile*, *Blackcurrant leaf*, *Devil's claw*.
21. A feed comprising the nutraceutical according to any of Claims 1-15.
22. A feed comprising the nutraceutical according to any of Claims 1-15 and grass meal.
23. A method of treating an animal suffering from an inflammatory condition or disease comprising administering to said animal an effective amount of at least one long chain polyunsaturated fatty acid and at least one carotenoid.
24. A method according to Claim 23 wherein said long chain fatty acid is a free fatty acid, or an ester thereof.
25. A method according to Claim 23 or 24 wherein said long chain fatty acid is selected from the group consisting of: a triglyceride, diglyceride, monoglyceride, phospholipids, glycolipid, sphingolipid or sulpholipid.

26. A method according to any of Claims 23-25 wherein said long chain fatty acid is docosahexanoic acid.
27. A method according to any of Claims 23-26 wherein said carotenoid is selected from the group consisting of: astaxanthin, zeaxanthin, lycopene, lutein, or carotene.
28. A method according to any of Claims 23-27 wherein said animal is administered a further anti-inflammatory agent.
29. A method according to any of Claims 23-28 wherein said disease or condition is selected from the group consisting of: inflammatory joint diseases e.g. rheumatoid arthritis, osteoarthritis, polyarthritis and gout; chronic inflammatory connective tissue diseases e.g. lupus erythematosus, scleroderma, Sjorgen's syndrome, poly- and dermatomyositis, vasculitis; mixed connective tissue disease (MCTD), e.g. tendonitis, synovitis, bacterial endocarditis, osteomyelitis and psoriasis; chronic inflammatory lung diseases e.g. chronic respiratory disease, pneumonia, fibrosing alveolitis, chronic bronchitis, bronchiectasis, emphysema, silicosis and other pneumoconiosis and tuberculosis; chronic inflammatory bowel and gastro-intestinal tract inflammatory diseases e.g. ulcerative colitis and Crohn's disease; chronic neural inflammatory diseases e.g. chronic inflammatory demyelinating polyradiculoneuropathy, chronic inflammatory demyelinating polyneuropathy, multiple sclerosis, Guillan-Barre Syndrome and myasthenia gravis; and including, mastitis, laminitis, laryngitis, chronic cholecystitis, Hashimoto's thyroiditis, inflammatory breast disease; chronic inflammation (e.g., caused by an implanted foreign body in a wound); and acute inflammatory tissue damage due to muscle damage after eccentric exercise (e.g., delayed onset muscle soreness - DOMS).
30. A method according to any of Claims 23-29 wherein said animal is selected from the group consisting of: horse; dog; cat ; cow; sheep; goat; mink; pig; hamster; mouse; rabbit; pot bellied pig; rat, gerbil, guinea pig.
31. A method according to any of Claims 23-29 wherein said animal is a human.

**Abstract**

5 We describe a composition comprising a source of long chain polyunsaturated fatty acid, for example, docosahexaenoic acid (DHA), and a carotenoid, for example, astaxanthin and other nutrients for prophylactic and/or therapeutic use in the healing of trauma- and stress-induced inflammatory conditions.



**PCT/GB2004/002707**



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ ~~FADED~~ TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ ~~GRAY~~ SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**